arXiv:1301.5401v2 [math.PR] 30 Jan 2013

Weingarten calculus for matrix ensembles associated with compact symmetric spaces

Sho Matsumoto

Abstract

A method for computing integrals of polynomial functions on compact symmetric spaces is given. Those integrals are expressed as sums of functions on symmetric groups.

1 Introduction

Let G be the unitary group $\mathrm{U}(n)$, orthogonal group $\mathrm{O}(n)$, or (compact) symplectic group $\mathrm{Sp}(2n)$, equipped with its Haar probability measure, and suppose that G is realized as a matrix group. Consider a random matrix $X = (x_{ij})$ picked up from G. Moments of matrix elements

$$\begin{aligned} \mathbb{E}[x_{i_1 j_1} \cdots x_{i_k j_k} \overline{x_{i'_1 j'_1} \cdots x_{i'_l j'_l}}] \qquad & \text{for } G = \mathrm{U}(n), \\ \mathbb{E}[x_{i_1 j_1} \cdots x_{i_k j_k}] \qquad & \text{for } G = \mathrm{O}(n) \text{ or } \mathrm{Sp}(2n) \end{aligned} \tag{1.1}$$

have been one of main interesting objects in random matrix theory. Weingarten [W] tried to understand the moments (1.1) for large n. Collins [Co] established a method of computations of (1.1) for $\mathrm{U}(n)$ with all finite n and called his method the *Weingarten calculus*. The moments are expressed as sums in terms of class functions $\mathrm{Wg}^{\mathrm{U}}(\cdot; n)$ on the symmetric group S_k. The function $\mathrm{Wg}^{\mathrm{U}}(\cdot; n)$, called the unitary Weingarten function, has rich combinatorial structures involving Jucys-Murphy elements [MN, N]. The Weingarten calculus for $\mathrm{O}(n)$ was constructed in [CSn, CM], and the corresponding Weingarten functions, called orthogonal Weingarten functions, are H_k-biinvariant functions on S_{2k}, where H_k is the hyperoctahedral subgroup of S_{2k}. The orthogonal Weingarten function also bears nice combinatorics [M1, Z]. The Weingarten calculus for $\mathrm{Sp}(2n)$ was mentioned in [CSn, CSt]. See also a recent preprint [Da], in which Brownian motions on classical groups are studied. In the last decade, Weingarten calculus has been widely applied: Harish-Chandra-Itzykson-Zuber integrals [Co, GGJ], quantum information theory [CN, CFN], designs [S], and statistics [CMS].

In his pioneering works for statistical mechanics, Dyson [Dy] introduced three important classes of random unitary matrices as modifications of Gaussian matrix ensembles.

These unitary matrix ensembles are known as *circular orthogonal/unitary/symplectic ensembles* (COE/CUE/CSE). Whereas the CUE is nothing but the unitary group equipped with the Haar measure, the COE and CSE are not Lie groups but compact symmetric spaces.

Generally, a classical compact symmetric space is of the form G/K, where G is a unitary, orthogonal, or symplectic group, and K is a closed subgroup of G consisting of fixed-points of an involution Ω on G. Let $\mathcal{S}$ be the image of the map $G \ni g \mapsto \Omega(g)^{-1}g$. We thus obtain a matrix realization of G/K:

$$G/K \simeq \mathcal{S}; \qquad G \ni g \mapsto \Omega(g)^{-1}g \in \mathcal{S}, \qquad K = \{h \in G \mid \Omega(h) = h\}.$$

Moreover, a G-invariant probability measure on $\mathcal{S}$ is induced from the Haar measure on G via this map. Then the matrix space $\mathcal{S}$ equipped with the probability measure is the matrix ensemble associated with G/K. In the case of the COE, $G/K = \mathrm{U}(n)/\mathrm{O}(n)$, $\Omega(g) = \overline{g}$ (the complex-conjugation of g), and $\mathcal{S}$ is the set of $n \times n$ symmetric unitary matrices. Cartan [Ca] classified classical compact symmetric spaces into seven infinite series labeled by A I, A II, A III, BD I, C I, C II, and D III (see Figure 1). Thus we obtain seven series of random matrix ensembles associated with these compact symmetric spaces. Eigenvalue distributions for them have been studied, see, e.g., [Du], [CaM], and [F, §3.7.2 and 3.7.3]. In contrast, distributions for those matrix elements were not studied so much. Ones can see a small work for only classes A I and A II in [F, §2.3.2].

In this article, we will construct Weingarten calculus for random matrices associated with each class $\mathcal{C} =$ A I, A II,... in Figure 1. A development in a similar direction can be seen in [CSt], but only asymptotic behaviors of two-degree moments were observed in that article. As in the case of classical groups $\mathrm{U}(n), \mathrm{O}(n), \mathrm{Sp}(2n)$, we will express the moments of random matrix elements as sums over symmetric groups. To do it, we need a distinguishing Weingarten function $\mathrm{Wg}^{\mathcal{C}}$ for each $\mathcal{C}$.

This article is organized as follows. In Section 2, we review the Weingarten calculus for $\mathrm{U}(n)$, $\mathrm{O}(n)$, and $\mathrm{Sp}(2n)$, developed in [Co, CM, CSn, CSt]. Especially, the case for symplectic groups is discussed in detail. It is described in [CSt, Da], however, unlike their descriptions, we introduce the symplectic Weingarten function in terms of twisted zonal spherical functions for the twisted Gelfand pair $(S_{2k}, H_k, \epsilon|_{H_k})$. In Section 3, we discuss the Weingarten calculus for random matrices from class A I and A II, which are the COE and CSE, respectively. The COE case ($\mathcal{C} =$ A I) is already given in [M2, M3] in detail but we present it here again for readers' convenience. In Section 4 and 5, we discuss the Weingarten calculus for chiral ensembles (A III, BD I, C II) and Bogolioubiov-de Gennes (BdG) ensembles (D III, C I), respectively. In the final section, we give a short conclusion for new Weingarten functions.

Class $\mathcal{C}$	Symmetric spaces	Random matrices
A I	$\mathrm{U}(n)/\mathrm{O}(n)$	circular orthogonal ensemble (COE)
A II	$\mathrm{U}(2n)/\mathrm{Sp}(2n)$	circular symplectic ensemble (CSE)
A III	$\mathrm{U}(n)/\mathrm{U}(a) \times \mathrm{U}(b) \quad (a+b=n)$	chiral ensemble
BD I	$\mathrm{O}(n)/\mathrm{O}(a) \times \mathrm{O}(b) \quad (a+b=n)$	
C II	$\mathrm{Sp}(2n)/\mathrm{Sp}(2a) \times \mathrm{Sp}(2b) \quad (a+b=n)$	
D III	$\mathrm{O}(2n)/\mathrm{U}(n)$	Bogolioubov-de Gennes (BdG) ensemble
C I	$\mathrm{Sp}(2n)/\mathrm{U}(n)$	

Figure 1: **List of classical compact symmetric spaces.** Remark that we consider the full groups $\mathrm{U}(n)$ and $\mathrm{O}(n)$ rather than special ones $\mathrm{SU}(n)$ and $\mathrm{SO}(n)$, the latter groups of which are usually used in Lie theory.

2 Weingarten calculus for classical groups

2.1 Weingarten calculus for unitary groups

In this subsection, we review the Weingarten calculus for unitary groups. See [Co, CSn, CMS] for details.

2.1.1 Partitions and cycle-types

A partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l)$ of a positive integer k is a weakly decreasing sequence of positive integers with $|\lambda| = \sum_{i=1}^{l} \lambda_i = k$. We write $\ell(\lambda)$ for the length l of λ. When λ is a partition of k, we write $\lambda \vdash k$.

Let S_k be the symmetric group acting on $[k] = \{1, 2, \dots, k\}$. A permutation $\sigma \in S_k$ is usually expressed in the two-array notation $\left(\begin{smallmatrix} 1 & 2 & \cdots & k \\ \sigma(1) & \sigma(2) & \cdots & \sigma(k) \end{smallmatrix}\right)$, but transpositions are often written as $(i\ j)$ shortly. Denote by id_k the identity permutation in S_k. If a permutation σ in S_k is decomposed into disjoint cycles with lengths $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_l$, then the partition $\mu = (\mu_1, \mu_2, \dots, \mu_l)$ of k is called the *cycle-type* of σ. For example, the cycle-type of the permutation $\left(\begin{smallmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 3 & 1 & 2 & 4 & 6 & 5 \end{smallmatrix}\right)$ in S_6 is $(3, 2, 1)$.

2.1.2 Unitary Weingarten functions

Let $L(S_k)$ be the algebra of complex-valued functions on S_k with convolution

$$(f_1 * f_2)(\sigma) = \sum_{\tau \in S_k} f_1(\tau) f_2(\tau^{-1}\sigma) \qquad (f_1, f_2 \in L(S_k),\ \sigma \in S_k).$$

The identity element in the algebra $L(S_k)$ is the Dirac function δ_{id_k}.

Let $\mathcal{Z}(L(S_k))$ be the center of $L(S_k)$: $\mathcal{Z}(L(S_k)) = \{h \in L(S_k) \mid h * f = f * h\ (f \in L(S_k))\}$. For a complex number z, we define the element $T^{\mathrm{U}}(\cdot; z)$ in $\mathcal{Z}(L(S_k))$ by

$$T^{\mathrm{U}}(\sigma; z) = z^{\ell(\mu)} \qquad (\sigma \in S_k),$$

where μ is the cycle-type of σ. Note that the upper index U stands for the unitary group. The class function $T^{\mathrm{U}}(\cdot;z)$ can be expanded in terms of irreducible characters χ^λ of S_k as follows:

$$T^{\mathrm{U}}(\cdot;z)=\frac{1}{k!}\sum_{\lambda\vdash k} f^\lambda C_\lambda(z)\chi^\lambda,$$

where $f^\lambda := \chi^\lambda(\mathrm{id}_k)$ is the dimension of the irreducible representation associated with λ and $C_\lambda(z)$ is the polynomial in z given by

$$C_\lambda(z)=\prod_{(i,j)\in\lambda}(z+j-i).$$

Here $(i,j)\in\lambda$ stands for $1\le i\le \ell(\lambda)$, $1\le j\le\lambda_i$. In other words, the (i,j) are coordinates of the Young diagram of λ.

The *unitary Weingarten function* $\mathrm{Wg}^{\mathrm{U}}(\cdot;z)$ on S_k is defined by

$$\mathrm{Wg}^{\mathrm{U}}(\cdot;z)=\frac{1}{k!}\sum_{\substack{\lambda\vdash k\\ C_\lambda(z)\neq 0}}\frac{f^\lambda}{C_\lambda(z)}\chi^\lambda, \tag{2.1}$$

summed over all partitions λ of k with $C_\lambda(z)\neq 0$. It is the pseudo-inverse element of $T^{\mathrm{U}}(\cdot;z)$, i.e., the unique element in $\mathcal{Z}(L(S_k))$ satisfying

$$T^{\mathrm{U}}(\cdot;z)*\mathrm{Wg}^{\mathrm{U}}(\cdot;z)*T^{\mathrm{U}}(\cdot;z)=T^{\mathrm{U}}(\cdot;z).$$

In particular, unless $z\in\{0,\pm1,\pm2,\dots,\pm(k-1)\}$, functions $T^{\mathrm{U}}(\cdot;z)$ and $\mathrm{Wg}^{\mathrm{U}}(\cdot;z)$ are inverse of each other and satisfy $T^{\mathrm{U}}(\cdot;z)*\mathrm{Wg}^{\mathrm{U}}(\cdot;z)=\delta_{\mathrm{id}_k}$.

2.1.3 Integrals on unitary groups

The unitary group $\mathrm{U}(n)$ is $\mathrm{U}(n)=\{U\in\mathrm{GL}(n,\mathbb{C})\mid UU^*=I_n\}$, which has the Haar probability measure. Here $U^*:=\overline{U^{\mathrm{T}}}$ is the adjoint matrix of U and $I_n=(\delta_{ij})_{1\le i,j\le n}$ is the $n\times n$ identity matrix. We will also write U for a random element of $\mathrm{U}(n)$, with distribution given by Haar measure.

Theorem 2.1 (Weingarten calculus for unitary groups [Co, CSn])**.** *Let $U=(u_{ij})_{1\le i,j\le n}$ be an $n\times n$ Haar unitary matrix. For four sequences $\boldsymbol{i}=(i_1,\dots,i_k)$, $\boldsymbol{j}=(j_1,\dots,j_k)$, $\boldsymbol{i'}=(i'_1,\dots,i'_k)$, $\boldsymbol{j'}=(j'_1,\dots,j'_k)$ of positive integers in $[n]$, we have*

$$\mathbb{E}[u_{i_1j_1}\cdots u_{i_kj_k}\overline{u_{i'_1j'_1}\cdots u_{i'_kj'_k}}]=\sum_{\sigma,\tau\in S_k}\delta_\sigma(\boldsymbol{i},\boldsymbol{i'})\delta_\tau(\boldsymbol{j},\boldsymbol{j'})\mathrm{Wg}^{\mathrm{U}}(\sigma^{-1}\tau;n).$$

Here $\delta_\sigma(\boldsymbol{i},\boldsymbol{i'})$ is defined by

$$\delta_\sigma(\boldsymbol{i},\boldsymbol{i'})=\prod_{s=1}^{k}\delta_{i_{\sigma(s)},i'_s}. \tag{2.2}$$

If $k \neq l$, *then* $\mathbb{E}[u_{i_1 j_1} \cdots u_{i_k j_k} \overline{u_{i'_1 j'_1} \cdots u_{i'_l j'_l}}]$ *vanishes for any indices* $i_1, \dots, i_k$, $j_1, \dots, j_k$, $i'_1, \dots, i'_l$, $j'_1, \dots, j'_l$.

Example 2.1. $\mathrm{Wg}^{\mathrm{U}}(\mathrm{id}_1; n) = \frac{1}{n}$;

$$\mathrm{Wg}^{\mathrm{U}}(\mathrm{id}_2; n) = \frac{1}{(n+1)(n-1)}; \qquad \mathrm{Wg}^{\mathrm{U}}((1\ 2); n) = \frac{-1}{n(n+1)(n-1)}.$$

We can see more examples in [Co].

Remark 2.1. In the sum of (2.1), we can drop the restriction for λ. Even if we replace the definition by

$$\mathrm{Wg}^{\mathrm{U}}(\cdot; z) = \frac{1}{k!} \sum_{\lambda \vdash k} \frac{f^\lambda}{C_\lambda(z)} \chi^\lambda,$$

which is a rational function in $z \in \mathbb{C}$ with finitely many poles, then Theorem 2.1 remains true after cancellations of poles. This enables us to ignore the restriction $C_\lambda(z) \neq 0$ in latter sections. See [CSn, Proposition 2.5] for details.

2.2 Weingarten calculus for orthogonal groups

In this subsection, we review the Weingarten calculus for orthogonal groups, developed in [CSn, CM, CMS]. The theory of zonal spherical functions for finite Gelfand pairs are seen in [Mac, Chapter VII].

2.2.1 Hyperoctahedral groups and coset-types

Let H_k be the hyperoctahedral subgroup of S_{2k} with order $2^k k!$, generated by adjacent transpositions $(2i-1\ 2i)$ $(1 \leq i \leq k)$ and double transpositions $(2i-1\ 2j-1)(2i\ 2j)$ $(1 \leq i < j \leq k)$. Let M_{2k} be the subset of permutations $\sigma \in S_{2k}$ satisfying

$$\sigma(2i-1) < \sigma(2i)\ (1 \leq i \leq k) \qquad \text{and} \qquad \sigma(1) < \sigma(3) < \cdots < \sigma(2k-1).$$

The elements σ in M_{2k} form the complete set of the representatives of S_{2k}/H_k and are sometimes identified with perfect matchings of the forms

$$\{\{\sigma(1), \sigma(2)\}, \{\sigma(3), \sigma(4)\}, \dots, \{\sigma(2k-1), \sigma(2k)\}\}$$

on $[2k]$.

For a permutation $\sigma \in S_{2k}$, we define the graph $\Gamma(\sigma)$ as follows. The vertex set is $\{1, 2, \dots, 2k\}$ and the edge set consists of red edges $\{2r-1, 2r\}$ and blue edges $\{\sigma(2r-1), \sigma(2r)\}$, where r runs over $1, 2, \dots, k$. Then all connected components of the graph $\Gamma(\sigma)$ have even vertices with numbers $2\mu_1 \geq 2\mu_2 \geq \cdots \geq 2\mu_l$. We call the partition $\mu := (\mu_1, \mu_2, \dots, \mu_l) \vdash k$ the *coset-type* of σ. For example, the coset-type of the permutation

$\left(\begin{smallmatrix}1&2&3&4&5&6\\3&1&2&4&6&5\end{smallmatrix}\right)$ in S_6 is $(2,1)$. The coset-type distinguishes with double cosets of H_k in S_{2k}. Specifically, for two permutations σ, τ in S_{2k}, their coset-types coincides if and only if $H_k \sigma H_k = H_k \tau H_k$ ([Mac, VII, (2.1)]).

For a partition $\mu = (\mu_1, \mu_2, \dots, \mu_l)$ of k, we define the permutation σ_μ in S_{2k} as follows: For each $r = 1, 2, \dots, l$,

$$\begin{aligned}
\sigma_\mu\left(2\textstyle\sum_{i=1}^{r-1}\mu_i + 1\right) &= 2\textstyle\sum_{i=1}^{r-1}\mu_i + 1, \\
\sigma_\mu\left(2\textstyle\sum_{i=1}^{r-1}\mu_i + 2\right) &= 2\textstyle\sum_{i=1}^{r-1}\mu_i + 2\mu_r, \\
\sigma_\mu\left(2\textstyle\sum_{i=1}^{r-1}\mu_i + p\right) &= 2\textstyle\sum_{i=1}^{r-1}\mu_i + p - 1 \quad \text{for } p = 3, 4, \dots, 2\mu_r.
\end{aligned} \tag{2.3}$$

For example, $\sigma_{(3,1)} = \left(\begin{smallmatrix}1&2&3&4&5&6&7&8\\1&6&2&3&4&5&7&8\end{smallmatrix}\right)$. It is easy to see that the permutation σ_μ belongs to M_{2k} and has the coset-type μ and signature $+1$. In particular, $\sigma_{(1^k)} = \mathrm{id}_{2k}$, the identity permutation in S_{2k}. We often regard σ_μ as a typical permutation of coset-type μ.

2.2.2 Orthogonal Weingarten functions and Gelfand pairs

Let $L(S_{2k}, H_k)$ be the subspace of all H_k-biinvariant functions in $L(S_{2k})$:

$$L(S_{2k}, H_k) = \{ f \in L(S_{2k}) \mid f(\zeta\sigma) = f(\sigma\zeta) = f(\sigma) \ (\sigma \in S_{2k},\ \zeta \in H_k) \}.$$

It is well known that (S_{2k}, H_k) is a *Gelfand pair*, i.e., $L(S_{2k}, H_k)$ is a commutative algebra under the convolution ([Mac, VII, (1.1) and (2.2)]).

For two functions f_1, f_2 in $L(S_{2k})$, we define a new product $f_1 \star f_2$ by

$$(f_1 \star f_2)(\sigma) = \sum_{\tau \in M_{2k}} f_1(\tau) f_2(\tau^{-1}\sigma) \qquad (\sigma \in S_{2k}).$$

Example 2.2. For $f_1, f_2 \in L(S_4)$,

$$\begin{aligned}
(f_1 \star f_2)\left(\begin{smallmatrix}1&2&3&4\\1&2&3&4\end{smallmatrix}\right) =& f_1\left(\begin{smallmatrix}1&2&3&4\\1&2&3&4\end{smallmatrix}\right) f_2\left(\begin{smallmatrix}1&2&3&4\\1&2&3&4\end{smallmatrix}\right) + f_1\left(\begin{smallmatrix}1&2&3&4\\1&3&2&4\end{smallmatrix}\right) f_2\left(\begin{smallmatrix}1&2&3&4\\1&3&2&4\end{smallmatrix}\right) + f_1\left(\begin{smallmatrix}1&2&3&4\\1&4&2&3\end{smallmatrix}\right) f_2\left(\begin{smallmatrix}1&2&3&4\\1&3&4&2\end{smallmatrix}\right). \\
(f_1 \star f_2)\left(\begin{smallmatrix}1&2&3&4\\1&3&2&4\end{smallmatrix}\right) =& f_1\left(\begin{smallmatrix}1&2&3&4\\1&2&3&4\end{smallmatrix}\right) f_2\left(\begin{smallmatrix}1&2&3&4\\1&3&2&4\end{smallmatrix}\right) + f_1\left(\begin{smallmatrix}1&2&3&4\\1&3&2&4\end{smallmatrix}\right) f_2\left(\begin{smallmatrix}1&2&3&4\\1&2&3&4\end{smallmatrix}\right) + f_1\left(\begin{smallmatrix}1&2&3&4\\1&4&2&3\end{smallmatrix}\right) f_2\left(\begin{smallmatrix}1&2&3&4\\1&4&3&2\end{smallmatrix}\right). \\
(f_1 \star f_2)\left(\begin{smallmatrix}1&2&3&4\\1&4&2&3\end{smallmatrix}\right) =& f_1\left(\begin{smallmatrix}1&2&3&4\\1&2&3&4\end{smallmatrix}\right) f_2\left(\begin{smallmatrix}1&2&3&4\\1&4&2&3\end{smallmatrix}\right) + f_1\left(\begin{smallmatrix}1&2&3&4\\1&3&2&4\end{smallmatrix}\right) f_2\left(\begin{smallmatrix}1&2&3&4\\1&4&3&2\end{smallmatrix}\right) + f_1\left(\begin{smallmatrix}1&2&3&4\\1&4&2&3\end{smallmatrix}\right) f_2\left(\begin{smallmatrix}1&2&3&4\\1&2&3&4\end{smallmatrix}\right).
\end{aligned}$$

For $f_1, f_2 \in L(S_{2k}, H_k)$, we have $f_1 \star f_2 = (2^k k!)^{-1} f_1 * f_2 \in L(S_{2k}, H_k)$. In fact, since M_{2k} is the complete set of representatives of cosets σH_k in S_{2k} and since f_1, f_2 are H_k-biinvariant, we have

$$(f_1 * f_2)(\sigma) = \sum_{\tau \in M_{2k}} \sum_{\zeta \in H_k} f_1(\tau\zeta) f_2((\tau\zeta)^{-1}\sigma) = \sum_{\tau \in M_{2k}} \sum_{\zeta \in H_k} f_1(\tau) f_2(\tau^{-1}\sigma) = |H_k| (f_1 \star f_2)(\sigma).$$

Hence $L(S_{2k}, H_k)$ is a commutative algebra under the product $\star$ with the identity element

$$\mathbf{1}_k(\sigma) = \begin{cases} 1 & \text{if } \sigma \in H_k, \\ 0 & \text{if } \sigma \in S_{2k} \setminus H_k. \end{cases}$$

For each partition $\lambda = (\lambda_1, \lambda_2, \dots) \vdash k$, we define the *zonal spherical function* ω^λ of the Gelfand pair (S_{2k}, H_k) by

$$\omega^\lambda = (2^k k!)^{-1} \chi^{2\lambda} * \mathbf{1}_k$$

with $2\lambda = (2\lambda_1, 2\lambda_2, \dots)$. The ω^λ, $\lambda \vdash k$, form a linear basis of $L(S_{2k}, H_k)$ and satisfy the orthogonality relation

$$\omega^\lambda \star \omega^\mu = \delta_{\lambda\mu} \frac{(2k)!}{2^k k!} \frac{1}{f^{2\lambda}} \omega^\lambda \tag{2.4}$$

([Mac, VII, (1.4)]).

For a complex number z, we define the element $T^{\mathrm{O}}(\sigma; z)$ in $L(S_{2k}, H_k)$ by

$$T^{\mathrm{O}}(\sigma; z) = z^{\ell(\mu)} \qquad (\sigma \in S_{2k}),$$

where μ is the coset-type of σ. We emphasize that T^{O} is different from T^{U}. The function $T^{\mathrm{O}}(\cdot; z)$ is expanded in terms of ω^λ as follows ([CM, (4.5)]):

$$T^{\mathrm{O}}(\cdot; z) = \frac{2^k k!}{(2k)!} \sum_{\lambda \vdash k} f^{2\lambda} D_\lambda(z) \omega^\lambda,$$

where $D_\lambda(z)$ is the polynomial in z given by

$$D_\lambda(z) = \prod_{(i,j) \in \lambda} (z + 2j - i - 1).$$

The *orthogonal Weingarten function* $\mathrm{Wg}^{\mathrm{O}}(\cdot; z)$ on S_{2k} is defined by

$$\mathrm{Wg}^{\mathrm{O}}(\cdot; z) = \frac{2^k k!}{(2k)!} \sum_{\substack{\lambda \vdash k \\ D_\lambda(z) \neq 0}} \frac{f^{2\lambda}}{D_\lambda(z)} \omega^\lambda,$$

which is the pseudo-inverse element of $T^{\mathrm{O}}(\cdot; z)$, i.e., the unique element in $L(S_k, H_k)$ satisfying

$$T^{\mathrm{O}}(\cdot; z) \star \mathrm{Wg}^{\mathrm{O}}(\cdot; z) \star T^{\mathrm{O}}(\cdot; z) = T^{\mathrm{O}}(\cdot; z). \tag{2.5}$$

In particular, if $D_\lambda(z) \neq 0$ for all partitions λ of k, functions $T^{\mathrm{O}}(\cdot; z)$ and $\mathrm{Wg}^{\mathrm{O}}(\cdot; z)$ are their inverse of each other and satisfy $T^{\mathrm{O}}(\cdot; z) \star \mathrm{Wg}^{\mathrm{O}}(\cdot; z) = \mathbf{1}_k$. Those relations follow from (2.4) and expansions of $T^{\mathrm{O}}(\cdot; z)$ and $\mathrm{Wg}^{\mathrm{O}}(\cdot; z)$ in terms of ω^λ.

2.2.3 Integrals on orthogonal groups

The (real) orthogonal group is $\mathrm{O}(n) = \{R \in \mathrm{GL}(n, \mathbb{R}) \mid RR^{\mathrm{T}} = I_n\}$ and has the Haar probability measure.

Theorem 2.2 (Weingarten calculus for orthogonal groups [CSn, CM])**.** *Let* $R = (r_{ij})_{1 \le i,j \le n}$ *be an* $n \times n$ *Haar orthogonal matrix. For two sequences* $\boldsymbol{i} = (i_1, \dots, i_{2k})$ *and* $\boldsymbol{j} = (j_1, \dots, j_{2k})$ *of positive integers in* $[n]$*, we have*

$$\mathbb{E}[r_{i_1 j_1} \cdots r_{i_{2k} j_{2k}}] = \sum_{\sigma, \tau \in M_{2k}} \Delta_\sigma(\boldsymbol{i}) \Delta_\tau(\boldsymbol{j}) \mathrm{Wg}^{\mathrm{O}}(\sigma^{-1}\tau; n).$$

Here $\Delta_\sigma(\boldsymbol{i})$ *is defined by*

$$\Delta_\sigma(\boldsymbol{i}) = \prod_{s=1}^{k} \delta_{i_{\sigma(2s-1)}, i_{\sigma(2s)}}. \tag{2.6}$$

Furthermore, $\mathbb{E}[r_{i_1 j_1} \cdots r_{i_{2k+1} j_{2k+1}}] = 0$ *for any* $i_1, \dots, i_{2k+1}, j_1, \dots, j_{2k+1}$*.*

Example 2.3. $\mathrm{Wg}^{\mathrm{O}}(\mathrm{id}_2; n) = \frac{1}{n}$;

$$\mathrm{Wg}^{\mathrm{O}}(\mathrm{id}_4; n) = \frac{n+1}{n(n+2)(n-1)}, \qquad \mathrm{Wg}^{\mathrm{O}}(\sigma_{(2)}; n) = \frac{-1}{n(n+2)(n-1)}.$$

We can see more examples in [CSn, CM].

2.3 Weingarten calculus for symplectic groups

In this subsection, we give the Weingarten calculus for symplectic groups. It was given in [CSt, Da], however, unlike their descriptions, we employ the theory of twisted Gelfand pairs ([Mac, VII, Examples 1-10, 1-11, 2-6, and 2-7]).

2.3.1 Twisted Gelfand pairs

Let ϵ be the signature function on S_{2k} and consider the linear space

$$L^\epsilon(S_{2k}, H_k) = \{ f \in L(S_{2k}) \mid f(\zeta\sigma) = f(\sigma\zeta) = \epsilon(\zeta) f(\sigma) \quad (\sigma \in S_{2k},\ \zeta \in H_k) \}.$$

The space $L^\epsilon(S_{2k}, H_k)$ is closed under the convolution $*$, and it becomes a $\mathbb{C}$-algebra. Furthermore, it is known that $L^\epsilon(S_{2k}, H_k)$ is a commutative algebra, which means that the triple $(S_{2k}, H_k, \epsilon|_{H_k})$ is, by definition, a *twisted Gelfand pair* ([Mac, VII, Example 2-6]). It is immediate to see that, if $f_1, f_2 \in L^\epsilon(S_{2k}, H_k)$, then $f_1 \star f_2 = (2^k k!)^{-1} f_1 * f_2 \in L^\epsilon(S_{2k}, H_k)$. Thus, $L^\epsilon(S_{2k}, H_k)$ is a commutative algebra under the product $\star$ with the identity element

$$\mathbf{1}_k^\epsilon(\sigma) = \begin{cases} \epsilon(\sigma) & \text{if } \sigma \in H_k, \\ 0 & \text{if } \sigma \in S_{2k} \setminus H_k. \end{cases}$$

For each partition $\lambda = (\lambda_1, \lambda_2, \dots) \vdash k$, we define the *twisted spherical function* π^λ of the twisted Gelfand pair by

$$\pi^\lambda = (2^k k!)^{-1} \chi^{\lambda \cup \lambda} * \mathbf{1}_k^\epsilon$$

with $\lambda \cup \lambda = (\lambda_1, \lambda_1, \lambda_2, \lambda_2, \dots)$. The map

$$L(S_{2k}, H_k) \to L^{\epsilon}(S_{2k}, H_k) : f \mapsto f^{\epsilon}, \qquad f^{\epsilon}(\sigma) = \epsilon(\sigma) f(\sigma) \ (\sigma \in S_{2k})$$

defines a $\mathbb{C}$-algebra isomorphism. The twisted spherical function π^{λ} is the image of $\omega^{\lambda'}$ with $\pi^{\lambda}(\sigma_{\mu}) = \omega^{\lambda'}(\sigma_{\mu})$. Here $\lambda' = (\lambda'_1, \lambda'_2, \dots)$ is the conjugate partition of λ, which is characterized by $(i,j) \in \lambda \Leftrightarrow (j,i) \in \lambda'$. Hence, the $\{\pi^{\lambda} \mid \lambda \vdash k\}$ form a linear basis of $L^{\epsilon}(S_{2k}, H_k)$ and satisfy the orthogonality relation

$$\pi^{\lambda} \star \pi^{\mu} = \delta_{\lambda\mu} \frac{(2k)!}{2^k k!} \frac{1}{f^{\lambda \cup \lambda}} \pi^{\lambda}.$$

2.3.2 Symplectic Weingarten functions

Let z be a complex number and consider the function $T^{\mathrm{Sp}}(\cdot; z)$ in $L^{\epsilon}(S_{2k}, H_k)$ defined by

$$T^{\mathrm{Sp}}(\sigma; z) = (-1)^k \epsilon(\sigma)(-2z)^{\ell(\mu)} \qquad (\sigma \in S_{2k}),$$

where μ is the coset-type of σ. Let

$$D'_{\lambda}(z) = \prod_{(i,j) \in \lambda} (2z - 2i + j + 1).$$

The expansion of $T^{\mathrm{Sp}}(\sigma; z)$ in terms of linear basis π^{λ} is given as follows.

Lemma 2.3.

$$T^{\mathrm{Sp}}(\cdot; z) = \frac{2^k k!}{(2k)!} \sum_{\lambda \vdash k} f^{\lambda \cup \lambda} D'_{\lambda}(z) \pi^{\lambda}. \tag{2.7}$$

Proof. We first suppose that $z = n$, a positive integer. We employ symmetric function theory. Recall power-sum symmetric functions p_{μ} and twisted zonal functions Z'_{λ}. We only use the following properties for them here (see [Mac, VII, Example 2-7]). For partitions λ, μ of k,

$$p_{\mu}(1^n) = n^{\ell(\mu)}, \qquad Z'_{\lambda}(1^n) = D'_{\lambda}(n),$$
$$p_{\mu} = \frac{2^k k!}{(2k)!} (-1)^{k-\ell(\mu)} 2^{-\ell(\mu)} \sum_{\lambda \vdash k} f^{\lambda \cup \lambda} \pi^{\lambda}(\sigma_{\mu}) Z'_{\lambda}.$$

Here $(1^n) = (1, 1, \dots, 1)$ with n times. Hence, if μ is the coset-type of $\sigma \in S_{2k}$, then

$$T^{\mathrm{Sp}}(\sigma; n) = \epsilon(\sigma)(-1)^{k-\ell(\mu)} 2^{\ell(\mu)} p_{\mu}(1^n) = \epsilon(\sigma) \frac{2^k k!}{(2k)!} \sum_{\lambda \vdash k} f^{\lambda \cup \lambda} \pi^{\lambda}(\sigma_{\mu}) D'_{\lambda}(n),$$

which implies the desired formula for $z = n$.

Since the both sides on (2.7) are polynomials in z, the equalities at all positive integers $z = n$ implies the ones at all complex numbers z. □

The *symplectic Weingarten function* with parameter z is the function in $L^{\epsilon}(S_{2k}, H_k)$ defined by

$$\mathrm{Wg}^{\mathrm{Sp}}(\cdot; z) = \frac{2^k k!}{(2k)!} \sum_{\substack{\lambda \vdash k \\ D'_\lambda(z) \neq 0}} \frac{f^{\lambda \cup \lambda}}{D'_\lambda(z)} \pi^\lambda. \tag{2.8}$$

Note that $T^{\mathrm{Sp}}(\sigma; z) = (-1)^k \epsilon(\sigma) T^{\mathrm{O}}(\sigma; -2z)$ and $\mathrm{Wg}^{\mathrm{Sp}}(\sigma; z) = (-1)^k \epsilon(\sigma) \mathrm{Wg}^{\mathrm{O}}(\sigma; -2z)$. Equation (2.5) is equivalent to

$$T^{\mathrm{Sp}}(\cdot; z) \star \mathrm{Wg}^{\mathrm{Sp}}(\cdot; z) \star T^{\mathrm{Sp}}(\cdot; z) = T^{\mathrm{Sp}}(\cdot; z). \tag{2.9}$$

2.3.3 Integrals on symplectic groups

Consider the vector space $\mathbb{C}^{2n}$ of column vectors with standard basis $(e_1, e_2, \dots, e_{2n})$. Define the skew-symmetric bilinear form $\langle \cdot, \cdot \rangle$ on $\mathbb{C}^{2n}$ by

$$\langle v, w \rangle = v^{\mathrm{T}} J w \tag{2.10}$$

with

$$J = J_n := \begin{pmatrix} O_n & I_n \\ -I_n & O_n \end{pmatrix}.$$

Let $(e_1^\vee, e_2^\vee, \dots, e_{2n}^\vee)$ be the dual basis of $(e_1, \dots, e_{2n})$ with respect to $\langle \cdot, \cdot \rangle$:

$$\langle e_i^\vee, e_j \rangle = \delta_{ij} \qquad (i, j \in [2n]).$$

More specifically, $(e_1^\vee, \dots, e_n^\vee, e_{n+1}^\vee, \dots, e_{2n}^\vee) = (-e_{n+1}, \dots, -e_{2n}, e_1, \dots, e_n)$. For convenience, we use the following notation:

$$\langle i, j \rangle := \langle e_i, e_j \rangle = \langle e_i^\vee, e_j^\vee \rangle = \begin{cases} 1 & \text{if } 1 \le i \le n \text{ and } j = i + n, \\ -1 & \text{if } 1 \le j \le n \text{ and } i = j + n, \\ 0 & \text{otherwise.} \end{cases}$$

Note that

$$e_i = J^{\mathrm{T}} e_i^\vee = \sum_{p=1}^{2n} \langle i, p \rangle e_p^\vee, \qquad e_i^\vee = J e_i = \sum_{p=1}^{2n} \langle p, i \rangle e_p \qquad (i = 1, 2, \dots, 2n) \tag{2.11}$$

and $J = (\langle i, j \rangle)_{1 \le i, j \le 2n}$.

For a $2n \times 2n$ matrix X, we define the *dual* matrix of X by

$$X^{\mathrm{D}} := J X^{\mathrm{T}} J^{\mathrm{T}}.$$

Then $\langle v, Xw \rangle = \langle X^{\mathrm{D}} v, w \rangle$ for all $v, w \in \mathbb{C}^{2n}$.

We realize the (unitary) symplectic group $\mathrm{Sp}(2n)$ by

$$\mathrm{Sp}(2n) = \{S = (s_{ij})_{1 \le i,j \le 2n} \in \mathrm{U}(2n) \mid SS^{\mathrm{D}} = I_{2n}\}.$$

It is equipped with the Haar probability measure. The following theorem was first given in [CSt] without the explicit expression (2.8) for $\mathrm{Wg}^{\mathrm{Sp}}$.

Theorem 2.4. *Let $S = (s_{ij})_{1 \le i,j \le 2n}$ be a Haar symplectic matrix. For two sequences $\boldsymbol{i} = (i_1, \dots, i_{2k})$ and $\boldsymbol{j} = (j_1, \dots, j_{2k})$ of positive integers in $[2n]$, we have*

$$\mathbb{E}[s_{i_1 j_1} s_{i_2 j_2} \cdots s_{i_{2k}, j_{2k}}] = \sum_{\sigma, \tau \in M_{2k}} \Delta'_\sigma(\boldsymbol{i}) \Delta'_\tau(\boldsymbol{j}) \mathrm{Wg}^{\mathrm{Sp}}(\sigma^{-1}\tau; n).$$

Here the symbol $\Delta'_\sigma(\boldsymbol{i}) \in \{0, 1, -1\}$ is defined by

$$\Delta'_\sigma(\boldsymbol{i}) := \prod_{r=1}^{k} \langle i_{\sigma(2r-1)}, i_{\sigma(2r)} \rangle. \tag{2.12}$$

Furthermore, $\mathbb{E}[s_{i_1 j_1} \cdots s_{i_{2k+1} j_{2k+1}}] = 0$ for any $i_1, \dots, i_{2k+1}, j_1, \dots, j_{2k+1}$.

We postpone the proof of this theorem in the next subsubsection.

Example 2.4. $\mathrm{Wg}^{\mathrm{Sp}}(\sigma; n) = \frac{\epsilon(\sigma)}{2n}$ for $\sigma \in S_2$;

$$\begin{aligned} \mathrm{Wg}^{\mathrm{Sp}}(\sigma; n) =& \epsilon(\sigma) \frac{2n-1}{4n(n-1)(2n+1)} && \text{for } \sigma \in H_2; \\ \mathrm{Wg}^{\mathrm{Sp}}(\sigma; n) =& \epsilon(\sigma) \frac{1}{4n(n-1)(2n+1)} && \text{for } \sigma \in S_4 \setminus H_2. \end{aligned}$$

2.3.4 Proof of Theorem 2.4

This proof was given by Collins and Stolz [CSt] (see also [CM]), however, they did not give any explicit expression for the symplectic Weingarten function. We here reconstruct their proof and observe how $\mathrm{Wg}^{\mathrm{Sp}}$ arises.

We first introduce the following useful notation. For a permutation $\sigma \in S_{2k}$ and two matrices $X = (x_{ij})_{1 \le i,j \le n}$ and $Y = (y_{ij})_{1 \le i,j \le n}$, put

$$\mathcal{T}_\sigma(X, Y) = \sum_{p_1, \dots, p_{2k} \in [n]} \prod_{r=1}^{k} x_{p_{\sigma(2r-1)}, p_{\sigma(2r)}} y_{p_{2r-1}, p_{2r}} = \sum_{q_1, \dots, q_{2k} \in [n]} \prod_{r=1}^{k} x_{q_{2r-1}, q_{2r}} y_{q_{\sigma^{-1}(2r-1)}, q_{\sigma^{-1}(2r)}}.$$

Note that $\mathcal{T}_{\sigma^{-1}}(X, Y) = \mathcal{T}_\sigma(Y, X)$ and $\mathcal{T}_{\sigma_\mu}(X, Y) = \prod_{i=1}^{\ell(\mu)} \mathcal{T}_{\sigma_{(\mu_i)}}(X, Y)$, where σ_μ is defined in (2.3). Since

$$\mathcal{T}_{\sigma_{(m)}}(X, Y) = \sum_{p_1, \dots, p_{2m}} x_{p_1, p_{2m}} y_{p_1, p_2} x_{p_2, p_3} y_{p_3, p_4} \cdots x_{p_{2m-2}, p_{2m-1}} \cdots y_{p_{2m-1}, p_{2m}},$$

we have

$$\mathcal{T}_{\sigma_{(m)}}(X,Y) = \begin{cases} \operatorname{Tr}[(XY)^m] & \text{if } X \text{ is a symmetric matrix,} \\ -\operatorname{Tr}[(XY)^m] & \text{if } X \text{ is a skew-symmetric matrix.} \end{cases} \tag{2.13}$$

As a particular case, we can see

$$\mathcal{T}_{\sigma}(J,J) = T^{\mathrm{Sp}}(\sigma;n) \qquad (\sigma \in S_{2k}). \tag{2.14}$$

Indeed, the skew-symmetry of J implies that the function $\sigma \mapsto \mathcal{T}_{\sigma}(J,J)$ belongs to $L^{\epsilon}(S_{2k}, H_k)$, and hence it is enough to check $\mathcal{T}_{\sigma_\mu}(J,J) = T^{\mathrm{Sp}}(\sigma_\mu;n)$ for all partitions μ. However, it follows from (2.13) that $\mathcal{T}_{\sigma_\mu}(J,J) = \prod_{i=1}^{\ell(\mu)} \mathcal{T}_{\sigma_{(\mu_i)}}(J,J) = \prod_{i=1}^{\ell(\mu)}[-\operatorname{Tr}(-I_{2n})^{\mu_i}] = (-1)^{k-\ell(\mu)}(2n)^{\ell(\mu)} = T^{\mathrm{Sp}}(\sigma_\mu;n)$.

We next recall the invariant theory for symplectic groups. Consider the tensor product $(\mathbb{C}^{2n})^{\otimes 2k}$ and define a bilinear form on $(\mathbb{C}^{2n})^{\otimes 2k}$ by

$$\left\langle \bigotimes_{j=1}^{2k} v_j, \bigotimes_{j=1}^{2k} w_j \right\rangle := \prod_{j=1}^{2k} \langle v_j, w_j \rangle, \qquad (v_1, \dots, v_{2k}, w_1, \dots, w_{2k} \in \mathbb{C}^{2n}),$$

where the skew-symmetric bilinear form $\langle v, w \rangle$ on the right hand side is defined in (2.10). Note that this bilinear form on $(\mathbb{C}^{2n})^{\otimes 2k}$ is symmetric.

Put

$$\theta_k = \sum_{p_1,\dots,p_k \in [2n]} e_{p_1}^{\vee} \otimes e_{p_1} \otimes \cdots \otimes e_{p_k}^{\vee} \otimes e_{p_k} \in (\mathbb{C}^{2n})^{\otimes 2k}.$$

The symmetric group S_{2k} acts on $(\mathbb{C}^{2n})^{\otimes 2k}$ by

$$\rho_{2k}(\sigma)(v_1 \otimes \cdots \otimes v_{2k}) = v_{\sigma^{-1}(1)} \otimes \cdots \otimes v_{\sigma^{-1}(2k)},$$

while the symplectic group $\mathrm{Sp}(2n)$ acts by

$$S(v_1 \otimes \cdots \otimes v_{2k}) = Sv_1 \otimes \cdots Sv_{2k}.$$

Then the First Fundamental Theorem for the symplectic group states that $\{\rho_{2k}(\sigma)\theta_k \mid \sigma \in M_{2k}\}$ spans the vector subspace of $(\mathbb{C}^{2n})^{\otimes 2k}$ consisting of invariant elements under the action of $\mathrm{Sp}(2n)$ (see, e.g., [GW, Theorem 5.3.4]).

Let us go back to the proof of Theorem 2.4. Let G be the symmetric matrix

$$\mathsf{G} = (\langle \rho_{2k}(\sigma)\theta_k, \rho_{2k}(\tau)\theta_k \rangle)_{\sigma,\tau \in M_{2k}} \tag{2.15}$$

and $\mathsf{W} = (\mathsf{w}(\sigma,\tau))_{\sigma,\tau \in M_{2k}}$ the pseudo-inverse matrix of G, i.e., W is the unique symmetric matrix satisfying $\mathsf{GWG} = \mathsf{G}$.

Let $S = (s_{ij})_{1 \le i,j \le 2n}$ be a Haar symplectic matrix. Since each matrix element s_{ij} is expressed as $s_{ij} = \langle e_i^\vee, S e_j \rangle$, we have

$$\begin{aligned}\mathbb{E}[s_{i_1 j_1} \cdots s_{i_{2k} j_{2k}}] =& \mathbb{E}\Big[\langle e_{i_1}^\vee \otimes \cdots \otimes e_{i_{2k}}^\vee, S(e_{j_1} \otimes \cdots \otimes e_{j_{2k}})\rangle\Big] \\ =& \Big\langle e_{i_1}^\vee \otimes \cdots \otimes e_{i_{2k}}^\vee, \mathbb{E}[S(e_{j_1} \otimes \cdots \otimes e_{j_{2k}})]\Big\rangle.\end{aligned}$$

As mentioned above, for any $\boldsymbol{v} \in V^{\otimes 2k}$, the $\mathrm{Sp}(2n)$-invariant vector $\mathbb{E}[S\boldsymbol{v}]$ can be expanded in terms of $\rho_{2k}(\sigma)\theta_k$. From a discussion parallel to the proof of [CM, Theorem 2.1] (see also [CSt]), the expansion is given by

$$\mathbb{E}[S\boldsymbol{v}] = \sum_{\sigma \in M_{2k}} \left(\sum_{\tau \in M_{2k}} \mathsf{w}(\sigma, \tau) \langle \boldsymbol{v}, \rho_{2k}(\tau)\theta_k \rangle \right) \rho_{2k}(\sigma)\theta_k.$$

Applying this to the previous equation, we obtain

$$\mathbb{E}[s_{i_1 j_1} \cdots s_{i_{2k} j_{2k}}] = \sum_{\sigma, \tau \in M_{2k}} \mathsf{w}(\sigma, \tau) \cdot \langle e_{i_1}^\vee \otimes \cdots \otimes e_{i_{2k}}^\vee, \rho_{2k}(\sigma)\theta_k \rangle \cdot \langle e_{j_1} \otimes \cdots \otimes e_{j_{2k}}, \rho_{2k}(\tau)\theta_k \rangle.$$

Here the values of bilinear forms are computed as follows:

$$\begin{aligned}&\langle e_{i_1}^\vee \otimes \cdots \otimes e_{i_{2k}}^\vee, \rho_{2k}(\sigma)\theta_k \rangle = \langle \rho_{2k}(\sigma^{-1})(e_{i_1}^\vee \otimes \cdots \otimes e_{i_{2k}}^\vee), \theta_k \rangle \\ =& \langle e_{i_{\sigma(1)}}^\vee \otimes \cdots \otimes e_{i_{\sigma(2k)}}^\vee, \theta_k \rangle = \prod_{r=1}^{k} \left(\sum_{p=1}^{2n} \langle e_{i_{\sigma(2r-1)}}^\vee, e_p^\vee \rangle \langle e_{i_{\sigma(2r)}}^\vee, e_p \rangle \right) \\ =& \Delta'_\sigma(\boldsymbol{i}),\end{aligned}$$

and, similarly, $\langle e_{j_1} \otimes \cdots \otimes e_{j_{2k}}, \rho_{2k}(\tau)\theta_k \rangle = \Delta'_\tau(\boldsymbol{j})$. In conclusion, Theorem 2.4 follows from the next lemma.

Lemma 2.5. *For* $\sigma, \tau \in M_{2k}$, $\mathsf{w}(\sigma, \tau) = \mathrm{Wg}^{\mathrm{Sp}}(\sigma^{-1}\tau; n)$.

Proof. If we put

$$\widetilde{T}(\sigma) = \langle \theta_k, \rho_{2k}(\sigma)\theta_k \rangle \qquad (\sigma \in S_{2k}),$$

then $\mathsf{G} = (\widetilde{T}(\sigma^{-1}\tau))_{\sigma, \tau \in M_{2k}}$. Using (2.11) we have

$$\begin{aligned}\theta_k =& \sum_{p_1, \dots, p_{2k}} \langle p_1, p_2 \rangle \langle p_3, p_4 \rangle \cdots \langle p_{2k-1}, p_{2k} \rangle e_{p_1}^\vee \otimes e_{p_2}^\vee \otimes \cdots \otimes e_{p_{2k}}^\vee \\ =& \sum_{q_1, \dots, q_{2k}} \langle q_1, q_2 \rangle \langle q_3, q_4 \rangle \cdots \langle q_{2k-1}, q_{2k} \rangle e_{q_1} \otimes e_{q_2} \otimes \cdots \otimes e_{q_{2k}},\end{aligned}$$

and hence

$$\tilde{T}(\sigma) = \sum_{q_1, \dots, q_{2k}} \langle q_{\sigma(1)}, q_{\sigma(2)} \rangle \cdots \langle q_{\sigma(2k-1)}, q_{\sigma(2k)} \rangle \langle q_1, q_2 \rangle \cdots \langle q_{2k-1}, q_{2k} \rangle = \mathcal{T}_\sigma(J, J),$$

which implies $\widetilde{T}(\sigma) = T^{\mathrm{Sp}}(\sigma; n)$ by (2.14). The matrix $(\mathrm{Wg}^{\mathrm{Sp}}(\sigma^{-1}\tau; n))_{\sigma, \tau \in M_{2k}}$ is therefore the pseudo-inverse matrix of $\mathsf{G} = (T^{\mathrm{Sp}}(\sigma^{-1}\tau; n))_{\sigma, \tau \in M_{2k}}$ by (2.9). $\square$

3 Circular ensembles

From now we consider random matrix ensembles $\mathcal{S}$ associated with classical symmetric spaces G/K. As we mentioned in Introduction, such ensembles are realized in the following way.

$$G/K \simeq \mathcal{S}; \qquad G \ni g \mapsto \Omega(g)^{-1}g \in \mathcal{S}.$$

Here Ω is an involution on G and K is the fixed-point set of Ω. If X is a Haar random matrix picked up from G, then the matrix $V := \Omega(X)^{-1}X$ is a random matrix associated with G/K. We consider the seven series of random matrices associated with compact symmetric spaces given in Figure 1.

In this section, we deal with the most important classes: circular orthogonal ensembles (COE) and circular symplectic ensembles (CSE).

3.1 Class A I

The setting for A I: $G = \mathrm{U}(n)$, $K = \mathrm{O}(n)$, $\Omega(g) = \overline{g}$. $\mathcal{S}$ consists of $n \times n$ symmetric unitary matrices.

When U is an $n\times n$ Haar unitary matrix, a random matrix corresponding to $\mathrm{U}(n)/\mathrm{O}(n)$ is defined by

$$V = V^{\mathrm{A\,I}} = \Omega(U)^{-1}U = U^{\mathrm{T}}U$$

and is said to be a COE matrix. The Weingarten calculus for a COE matrix is constructed in [M2]. For completeness of this paper, we review main results in [M2]. Applying the Weingarten calculus for $\mathrm{U}(n)$, we can obtain the following theorem.

Theorem 3.1 (Theorem 1.1 and Proposition 3.1 in [M2]). *Let* $V = V^{\mathrm{A\,I}} = (v_{ij})_{1\le i,j\le n}$ *be an* $n \times n$ *COE matrix. For two sequences* $\boldsymbol{i} = (i_1, \dots, i_{2k})$ *and* $\boldsymbol{j} = (j_1, \dots, j_{2k})$*, we have*

$$\mathbb{E}[v_{i_1i_2}v_{i_3i_4}\cdots v_{i_{2k-1}i_{2k}}\overline{v_{j_1j_2}v_{j_3j_4}\cdots v_{j_{2k-1}j_{2k}}}] = \sum_{\sigma\in S_{2k}} \delta_\sigma(\boldsymbol{i},\boldsymbol{j})\mathrm{Wg}^{\mathrm{A\,I}}(\sigma;n)$$

with the convolution $\mathrm{Wg}^{\mathrm{A\,I}}(\cdot;n) := T^{\mathrm{O}}(\cdot;n) * \mathrm{Wg}^{\mathrm{U}}(\cdot;n)$ *in* $L(S_{2k})$*. If* $k \neq l$ *then*

$$\mathbb{E}[v_{i_1i_2}v_{i_3i_4}\cdots v_{i_{2k-1}i_{2k}}\overline{v_{j_1j_2}v_{j_3j_4}\cdots v_{j_{2l-1}j_{2l}}}]$$

always vanishes. Moreover, the function $\mathrm{Wg}^{\mathrm{A\,I}}(\cdot;n)$ *belongs to* $L(S_{2k}, H_k)$ *and coincides with the orthogonal Weingarten function with parameter* $n+1$*, i.e.,*

$$\mathrm{Wg}^{\mathrm{A\,I}}(\sigma;n) = \mathrm{Wg}^{\mathrm{O}}(\cdot;n+1).$$

As a corollary of this theorem, the following moments for a single entry are computed (see [M2, Theorems 4.1 and 4.2]):

$$\begin{aligned}\mathbb{E}[|v_{ii}|^{2k}] =& \frac{2^k k!}{(n+1)(n+3)\cdots(n+2k-1)},\\ \mathbb{E}[|v_{ij}|^{2k}] =& \frac{k!}{n(n+1)(n+2)\cdots(n+k-2)(n+2k-1)} \qquad (i \neq j).\end{aligned}$$

The first equation can be obtained easily from the previous theorem, but the derivation of the second one is somewhat complicated.

Example 3.1. From Example 2.3 we have $\mathrm{Wg}^{\mathrm{A\,I}}(\sigma;n) = \mathrm{Wg}^{\mathrm{O}}(\sigma;n+1) = \frac{1}{n+1}$ for $\sigma \in S_2$;

$$\mathrm{Wg}^{\mathrm{A\,I}}(\sigma;n) = \mathrm{Wg}^{\mathrm{O}}(\sigma;n+1) = \begin{cases} \frac{n+2}{n(n+1)(n+3)} & \text{for } \sigma \in H_2 \\ \frac{-1}{n(n+1)(n+3)} & \text{for } \sigma \in S_4 \setminus H_2. \end{cases}$$

3.2 Class A II

The setting for A II: $G = \mathrm{U}(2n)$, $K = \mathrm{Sp}(2n)$, $\Omega(g) = (g^{\mathrm{D}})^{-1}$. $\mathcal{S}$ consists of $2n \times 2n$ unitary matrices g satisfying $g^{\mathrm{D}} = g$.

When U is a $2n\times 2n$ Haar unitary matrix, a random matrix corresponding to $\mathrm{U}(2n)/\mathrm{Sp}(2n)$ is defined by

$$V = V^{\mathrm{A\,II}} = U^{\mathrm{D}} U$$

and is said to be a CSE matrix.

We would like to compute mixed moments for v_{ij} and $\overline{v_{ij}}$. In order to simplify the notation, we deal with

$$\tilde{v}_{ij} := \langle e_i, V e_j \rangle \qquad (1 \le i, j \le 2n)$$

instead of $v_{ij} = \langle e_i^{\vee}, V e_j \rangle$. More specifically,

$$v_{ij} = \begin{cases} -\tilde{v}_{i+n,j} & \text{if } 1 \le i \le n, \\ \tilde{v}_{i-n,j} & \text{if } n+1 \le i \le 2n. \end{cases}$$

Theorem 3.2. *Let V be a $2n \times 2n$ CSE matrix. For two sequences $\boldsymbol{i} = (i_1, \dots, i_{2k})$ and $\boldsymbol{j} = (j_1, \dots, j_{2k})$ in $[2n]^{\times 2k}$, we have*

$$\mathbb{E}[\tilde{v}_{i_1,i_2}\tilde{v}_{i_3,i_4}\cdots\tilde{v}_{i_{2k-1},i_{2k}}\overline{\tilde{v}_{j_1,j_2}\tilde{v}_{j_3,j_4}\cdots\tilde{v}_{j_{2k-1},j_{2k}}}] = \sum_{\sigma \in S_{2k}} \delta_\sigma(\boldsymbol{i},\boldsymbol{j}) \mathrm{Wg}^{\mathrm{A\,II}}(\sigma;n)$$

with $\mathrm{Wg}^{\mathrm{A\,II}}(\cdot;n) := T^{\mathrm{Sp}}(\cdot;n) * \mathrm{Wg}^{\mathrm{U}}(\cdot;2n)$. *If $k \neq l$ then*

$$\mathbb{E}[\tilde{v}_{i_1,i_2}\tilde{v}_{i_3,i_4}\cdots\tilde{v}_{i_{2k-1},i_{2k}}\overline{\tilde{v}_{j_1,j_2}\tilde{v}_{j_3,j_4}\cdots\tilde{v}_{j_{2l-1},j_{2l}}}] = 0.$$

Proof. Each element of a $2n \times 2n$ CSE matrix $V = U^{\mathrm{D}}U$ is given as

$$\tilde{v}_{i,i'} = \langle e_i, U^{\mathrm{D}}Ue_{i'}\rangle = \langle Ue_i, Ue_{i'}\rangle = \sum_{p,q\in[2n]} u_{pi}u_{qi'}\langle p,q\rangle \qquad (i,i' \in [2n]).$$

Hence

$$\begin{aligned}&\mathbb{E}[\tilde{v}_{i_1,i_2}\tilde{v}_{i_3,i_4}\cdots\tilde{v}_{i_{2k-1},i_{2k}}\overline{\tilde{v}_{j_1,j_2}\tilde{v}_{j_3,j_4}\cdots\tilde{v}_{j_{2k-1},j_{2k}}}]\\ =&\sum_{\boldsymbol{p}=(p_1,\dots,p_{2k})}\sum_{\boldsymbol{q}=(q_1,\dots,q_{2k})}\mathbb{E}[u_{p_1i_1}\cdots u_{p_ki_k}\overline{u_{q_1j_1}\cdots u_{q_kj_k}}]\prod_{r=1}^{k}\langle p_{2r-1},p_{2r}\rangle\langle q_{2r-1},q_{2r}\rangle.\end{aligned}$$

(When $k \neq l$, $\mathbb{E}[u_{p_1i_1}\cdots u_{p_ki_k}\overline{u_{q_1j_1}\cdots u_{q_lj_l}}] \equiv 0$.) Applying the Weingarten calculus for a Haar unitary matrix U, we have

$$\mathbb{E}[\tilde{v}_{i_1,i_2}\tilde{v}_{i_3,i_4}\cdots\tilde{v}_{i_{2k-1},i_{2k}}\overline{\tilde{v}_{j_1,j_2}\tilde{v}_{j_3,j_4}\cdots\tilde{v}_{j_{2k-1},j_{2k}}}] = \sum_{\sigma\in S_{2k}}\delta_\sigma(\boldsymbol{i},\boldsymbol{j})\sum_{\tau\in S_{2k}}\mathrm{Wg}^{\mathrm{U}}(\tau^{-1}\sigma;2n)\widetilde{T}(\tau),$$

where

$$\begin{aligned}\widetilde{T}(\tau) =&\sum_{\boldsymbol{p}=(p_1,\dots,p_{2k})\in[2n]^{\times 2k}}\sum_{\boldsymbol{q}=(q_1,\dots,q_{2k})\in[2n]^{\times 2k}}\delta_\tau(\boldsymbol{p},\boldsymbol{q})\prod_{r=1}^{k}\langle p_{2r-1},p_{2r}\rangle\langle q_{2r-1},q_{2r}\rangle\\ =&\mathcal{T}_\tau(J,J) = T^{\mathrm{Sp}}(\tau;n).\end{aligned}$$

The last equality above follows from (2.14). □

Proposition 3.3. *The function* $\mathrm{Wg}^{\mathrm{A\,II}}(\cdot;n) = T^{\mathrm{Sp}}(\cdot;n) * \mathrm{Wg}^{\mathrm{U}}(\cdot;2n)$ *coincides with the symplectic Weingarten function with parameter* $n-\frac{1}{2}$. *Specifically, for each* $\sigma \in S_{2k}$, *we have*

$$\mathrm{Wg}^{\mathrm{A\,II}}(\sigma;n) = \mathrm{Wg}^{\mathrm{Sp}}(\sigma;n-\tfrac{1}{2}).$$

Proof. It follows from (2.7) and (2.1) that

$$T^{\mathrm{Sp}}(\cdot;n) * \mathrm{Wg}^{\mathrm{U}}(\cdot;2n) = \frac{2^k k!}{(2k)!}\sum_{\lambda\vdash k} f^{\lambda\cup\lambda}D'_\lambda(n)\frac{1}{(2k)!}\sum_{\mu\vdash 2k}\frac{f^\mu}{C_\mu(2n)}\pi^\lambda * \chi^\mu.$$

Since $\pi^\lambda * \chi^\mu = \delta_{\lambda\cup\lambda,\mu}\frac{(2k)!}{f^{\lambda\cup\lambda}}\pi^\lambda$ and since

$$\frac{D'_\lambda(z)}{C_{\lambda\cup\lambda}(2z)} = \frac{\prod_{(i,j)\in\lambda}(2z-2i+j+1)}{\prod_{(i,j)\in\lambda}(2z+j-(2i-1))(2z+j-2i)} = \frac{1}{\prod_{(i,j)\in\lambda}(2z+j-2i)} = \frac{1}{D'_\lambda(z-\frac{1}{2})},$$

we have $T^{\mathrm{Sp}}(\cdot;n) * \mathrm{Wg}^{\mathrm{U}}(\cdot;2n) = \frac{2^k k!}{(2k)!}\sum_{\lambda\vdash k}\frac{f^{\lambda\cup\lambda}}{D'_\lambda(n-\frac{1}{2})}\pi^\lambda = \mathrm{Wg}^{\mathrm{Sp}}(\cdot;n-\frac{1}{2})$. □

Example 3.2. From Example 2.4 we have $\mathrm{Wg}^{\mathrm{A\,II}}(\sigma;n) = \mathrm{Wg}^{\mathrm{Sp}}(\sigma;n-\frac{1}{2}) = \frac{\epsilon(\sigma)}{2n-1}$ for $\sigma \in S_2$;

$$\mathrm{Wg}^{\mathrm{A\,II}}(\sigma;n) = \mathrm{Wg}^{\mathrm{Sp}}(\sigma;n-\tfrac{1}{2}) = \begin{cases}\epsilon(\sigma)\frac{n-1}{n(2n-1)(2n-3)} & \text{for } \sigma \in H_2,\\ \epsilon(\sigma)\frac{1}{2n(2n-1)(2n-3)} & \text{for } \sigma \in S_4 \setminus H_2.\end{cases}$$

4 Chiral ensembles

In this section, we deal with random matrix ensembles associated with classes A III, BD I, C II. They are known as chiral ensembles.

4.1 Class A III

The setting for A III: $G = \mathrm{U}(n)$, $K = \mathrm{U}(a) \times \mathrm{U}(b)$, $\Omega(g) = I'_{ab} g I'_{ab}$, where $n = a + b$ with $a \geq b \geq 1$ and

$$I'_{ab} = \begin{pmatrix} I_a & O \\ O & -I_b \end{pmatrix}.$$

Let U be an $n \times n$ Haar unitary matrix. A random matrix corresponding to $\mathrm{U}(n)/\mathrm{U}(a) \times \mathrm{U}(b) = \mathrm{SU}(n)/\mathrm{S}(\mathrm{U}(a) \times \mathrm{U}(b))$ is defined by $V = V^{\mathrm{A\,III}} = I'_{ab} U^* I'_{ab} U$. For the sake of ease, we consider a Hermitian and unitary random matrix

$$W = W^{\mathrm{A\,III}} = U^* I'_{ab} U,$$

instead of $V = I'_{ab} W$.

We define the function $T^{\mathrm{A\,III}}_{ab}$ in $\mathscr{Z}(L(S_k))$ by

$$T^{\mathrm{A\,III}}_{ab}(\sigma) = \mathrm{Tr}_\sigma(I'_{ab})$$

where $\mathrm{Tr}_\sigma(A) = \prod_{j=1}^{\ell(\mu)} \mathrm{Tr}(A^{\mu_j})$ if μ is the cycle-type of σ. More specifically, we have

$$T^{\mathrm{A\,III}}_{ab}(\sigma) = (a+b)^{\ell^{\mathrm{e}}(\mu)} (a-b)^{\ell^{\mathrm{o}}(\mu)}.$$

Here $\ell^{\mathrm{e}}(\mu)$ (resp. $\ell^{\mathrm{o}}(\mu)$) is the number of parts μ_j with even lengths (resp. odd lengths).

Theorem 4.1. *Let $W = W^{\mathrm{A\,III}}$ be the random matrix defined as above. For two sequences $\boldsymbol{i} = (i_1, \dots, i_k)$ and $\boldsymbol{j} = (j_1, \dots, j_k)$, we have*

$$\mathbb{E}[w_{i_1 j_1} w_{i_2 j_2} \cdots w_{i_k j_k}] = \sum_{\sigma \in S_k} \delta_\sigma(\boldsymbol{i}, \boldsymbol{j}) \mathrm{Wg}^{\mathrm{A\,III}}(\sigma; a, b).$$

Here the function $\mathrm{Wg}^{\mathrm{A\,III}}(\cdot; a, b)$ *in* $\mathscr{Z}(L(S_k))$ *is defined by*

$$\mathrm{Wg}^{\mathrm{A\,III}}(\cdot; a, b) = T^{\mathrm{A\,III}}_{ab} * \mathrm{Wg}^{\mathrm{U}}(\cdot; n).$$

Proof. The random matrix W is Hermitian and the distribution of W is invariant under the unitary transform $W \mapsto g W g^*$, where g is a fixed unitary matrix. Hence we can apply Theorem 3.1 in [CMS] and obtain

$$\mathbb{E}[w_{i_1 j_1} w_{i_2 j_2} \cdots w_{i_k j_k}] = \sum_{\sigma, \tau \in S_k} \delta_\sigma(\boldsymbol{i}, \boldsymbol{j}) \mathrm{Wg}^{\mathrm{U}}(\tau^{-1}\sigma; n) \mathbb{E}[\mathrm{Tr}_\tau(W)].$$

Here, wee see that $\mathbb{E}[\mathrm{Tr}_\tau(W)] = \mathbb{E}[\mathrm{Tr}_\tau(I'_{ab})] = \mathrm{Tr}_\tau(I'_{ab}) = T^{\mathrm{A\,III}}_{ab}(\tau)$, and we obtain the desired identity. ☐

Example 4.1. $n = a+b$. $\mathrm{Wg}^{\mathrm{A\,III}}(\mathrm{id}_1; a, b) = \frac{a-b}{n}$;

$$\mathrm{Wg}^{\mathrm{A\,III}}(\mathrm{id}_2; a, b) = \frac{(a-b+1)(a-b-1)}{(n+1)(n-1)}, \qquad \mathrm{Wg}^{\mathrm{A\,III}}((1\ 2); a, b) = \frac{4ab}{n(n-1)(n+1)}.$$

Recall power symmetric functions p_μ and Schur functions s_λ. They have the relation

$$p_\mu = \sum_{\lambda \vdash k} \chi^\lambda(\sigma) s_\lambda$$

if $\mu \vdash k$ is the cycle-type of σ ([Mac, I, (7.8)]). Furthermore, it is easy to see that

$$p_\mu(1^a, (-1)^b) = p_\mu(\underbrace{1, 1, \dots, 1}_{a}, \underbrace{-1, -1, \dots, -1}_{b}) = T^{\mathrm{A\,III}}_{ab}(\sigma).$$

Hence we obtain the expansion of $T^{\mathrm{A\,III}}_{ab}$ in terms of irreducible characters χ^λ:

$$T^{\mathrm{A\,III}}_{ab} = \sum_{\lambda \vdash k} s_\lambda(1^a, (-1)^b) \chi^\lambda.$$

On the other hand, from the well-known identity $s_\lambda(1^n) = \frac{f^\lambda C_\lambda(n)}{k!}$, the unitary Weingarten function is expressed as

$$\mathrm{Wg}^{\mathrm{U}}(\cdot; n) = \frac{1}{(k!)^2} \sum_{\lambda \vdash k} \frac{(f^\lambda)^2}{s_\lambda(1^n)} \chi^\lambda.$$

Consequently, using the relation $\chi^\lambda * \chi^\mu = \frac{k!}{f^\lambda} \delta_{\lambda\mu} \chi^\lambda$, we obtain the following expansion of $\mathrm{Wg}^{\mathrm{A\,III}}(\cdot; a, b)$ in terms of χ^λ:

$$\mathrm{Wg}^{\mathrm{A\,III}}(\cdot; a, b) = T^{\mathrm{A\,III}}_{ab} * \mathrm{Wg}^{\mathrm{U}}(\cdot; n) = \frac{1}{k!} \sum_{\lambda \vdash k} f^\lambda \frac{s_\lambda(1^a, (-1)^b)}{s_\lambda(1^{a+b})} \chi^\lambda.$$

4.2 Class BD I

The setting for BD I: $G = \mathrm{O}(n)$, $K = \mathrm{O}(a) \times \mathrm{O}(b)$, $\Omega(g) = I'_{ab} g I'_{ab}$, where $n = a+b$ with $a \geq b \geq 1$.

The discussion is parallel to that in the previous subsection. We deal with the symmetric and orthogonal random matrix

$$W = W^{\mathrm{BD\,I}} = R^{\mathrm{T}} I'_{ab} R,$$

where R is an $n \times n$ Haar orthogonal matrix. This random matrix is associated with the symmetric space $\mathrm{O}(n)/\mathrm{O}(a) \times \mathrm{O}(b) = \mathrm{SO}(n)/\mathrm{S}(\mathrm{O}(a) \times \mathrm{O}(b))$.

We define the function $T^{\mathrm{BD\,I}}_{ab}$ in $L(S_{2k}, H_k)$ by

$$T^{\mathrm{BD\,I}}(\sigma) = \mathcal{T}_\sigma(I'_{ab}, I_n).$$

More specifically, if $\sigma \in S_{2k}$ has the coset-type $\mu = (\mu_1, \mu_2, \dots, \mu_l)$, then

$$T_{ab}^{\mathrm{BD\,I}}(\sigma) = \prod_{i=1}^{\ell(\mu)} \mathrm{Tr}(I'_{ab})^{\mu_i} = (a+b)^{\ell^{\mathrm{e}}(\mu)}(a-b)^{\ell^{\mathrm{o}}(\mu)}.$$

Theorem 4.2. *Let* $W = W^{\mathrm{BD\,I}}$ *be the random matrix defined as above. For a sequence* $\boldsymbol{i} = (i_1, \dots, i_{2k})$*, we have*

$$\mathbb{E}[w_{i_1 i_2} w_{i_3 i_4} \cdots w_{i_{2k-1} i_{2k}}] = \sum_{\sigma \in M_{2k}} \Delta_\sigma(\boldsymbol{i}) \mathrm{Wg}^{\mathrm{BD\,I}}(\sigma; a, b).$$

Here the function $\mathrm{Wg}^{\mathrm{BD\,I}}(\cdot; a, b)$ *in* $L(S_{2k}, H_k)$ *is defined by*

$$\mathrm{Wg}^{\mathrm{BD\,I}}(\cdot; a, b) = T_{ab}^{\mathrm{BD\,I}} \star \mathrm{Wg}^{\mathrm{O}}(\cdot; n).$$

Proof. The proof is the same with that of Theorem 4.1. Apply Theorem 3.3 in [CMS]. □

Example 4.2. $n = a + b$. $\mathrm{Wg}^{\mathrm{BD\,I}}(\mathrm{id}_2; a, b) = \frac{a-b}{n}$;

$$\begin{aligned}
\mathrm{Wg}^{\mathrm{BD\,I}}\left(\left(\begin{smallmatrix} 1&2&3&4 \\ 1&2&3&4 \end{smallmatrix}\right); a, b\right) &= \frac{(a-b)^2(n+1) - 2n}{n(n+2)(n-1)}, \\
\mathrm{Wg}^{\mathrm{BD\,I}}\left(\left(\begin{smallmatrix} 1&2&3&4 \\ 1&3&2&4 \end{smallmatrix}\right); a, b\right) = \mathrm{Wg}^{\mathrm{BD\,I}}\left(\left(\begin{smallmatrix} 1&2&3&4 \\ 1&4&2&3 \end{smallmatrix}\right); a, b\right) &= \frac{4ab}{n(n+2)(n-1)}.
\end{aligned}$$

Using zonal polynomials Z_λ (see [Mac, VII-2]), we can obtain the expansion of $\mathrm{Wg}^{\mathrm{BD\,I}}(\cdot; a, b)$ in terms of zonal spherical functions ω^λ:

$$\mathrm{Wg}^{\mathrm{BD\,I}}(\cdot; a, b) = \frac{2^k k!}{(2k)!} \sum_{\lambda \vdash k} f^{2\lambda} \frac{Z_\lambda(1^a, (-1)^b)}{Z_\lambda(1^{a+b})} \omega^\lambda.$$

4.3 Class C II

The setting for C II: $G = \mathrm{Sp}(2n)$, $K = \mathrm{Sp}(2a) \times \mathrm{Sp}(2b)$, $\Omega(g) = I''_{ab} g I''_{ab}$, where $n = a + b$ with $a \geq b \geq 1$ and

$$I''_{ab} = \begin{pmatrix} I'_{ab} & O \\ O & I'_{ab} \end{pmatrix}.$$

The discussion is parallel to the previous subsection again. We deal with the random matrix

$$W = W^{\mathrm{C\,II}} = S^{\mathrm{D}} I''_{ab} S.$$

where S is a $2n \times 2n$ Haar symplectic matrix. It is associated with the symmetric space $\mathrm{Sp}(2n)/\mathrm{Sp}(2a) \times \mathrm{Sp}(2b)$.

We define the function $T_{ab}^{\mathrm{C\,II}}$ in $L^{\epsilon}(S_{2k}, H_k)$ by

$$T_{ab}^{\mathrm{C\,II}}(\sigma) = \mathcal{T}_{\sigma}(J, JI''_{ab}).$$

If $\sigma \in S_{2k}$ has the coset-type μ, then

$$\begin{aligned} T_{ab}^{\mathrm{C\,II}}(\sigma) =& \epsilon(\sigma) \prod_{i=1}^{\ell(\mu)} T_{ab}^{\mathrm{C\,II}}(\sigma_{(\mu_i)}) = \epsilon(\sigma) \prod_{i=1}^{\ell(\mu)} [-\operatorname{Tr}(JJI''_{ab})^{\mu_i}] \\ =& (-1)^{k-\ell(\mu)} \epsilon(\sigma) \prod_{i=1}^{\ell(\mu)} \operatorname{Tr}(I''_{ab})^{\mu_i} = (-1)^{k-\ell(\mu)} \epsilon(\sigma) 2^{\ell(\mu)} (a+b)^{\ell^{\mathrm{e}}(\mu)} (a-b)^{\ell^{\mathrm{o}}(\mu)}. \end{aligned}$$

Here the second equality above follows by (2.13).

As in the case of class A II, we consider $\tilde{w}_{ij} = \langle e_i, W e_j \rangle$ instead of matrix elements w_{ij}.

Theorem 4.3. *Let $W = W^{\mathrm{C\,II}}$ be the random matrix defined as above. For a sequence $\boldsymbol{i} = (i_1, \dots, i_{2k})$, we have*

$$\mathbb{E}[\tilde{w}_{i_1,i_2} \tilde{w}_{i_3,i_4} \cdots \tilde{w}_{i_{2k-1},i_{2k}}] = \sum_{\sigma \in M_{2k}} \Delta'_{\sigma}(\boldsymbol{i}) \mathrm{Wg}^{\mathrm{C\,II}}(\sigma; a, b).$$

Here the function $\mathrm{Wg}^{\mathrm{C\,II}}(\cdot; a, b)$ *in* $L^{\epsilon}(S_{2k}, H_k)$ *is defined by*

$$\mathrm{Wg}^{\mathrm{C\,II}}(\cdot; a, b) = T_{ab}^{\mathrm{C\,II}} \star \mathrm{Wg}^{\mathrm{Sp}}(\cdot; n).$$

Proof. Since

$$\tilde{w}_{i,j} = \langle e_i, S^{\mathrm{D}} I''_{ab} S e_j \rangle = \langle S e_i, I''_{ab} S e_j \rangle = \sum_{p,q=1}^{2n} s_{pi} s_{qj} \langle e_p, I''_{ab} e_q \rangle,$$

we have

$$\mathbb{E}[\tilde{w}_{i_1,i_2} \tilde{w}_{i_3,i_4} \cdots \tilde{w}_{i_{2k-1},i_{2k}}] = \sum_{p_1,\dots,p_{2k}} \mathbb{E}[s_{p_1 i_1} s_{p_2 i_2} \cdots s_{p_{2k} i_{2k}}] \prod_{r=1}^{k} \langle p_{2r-1}, p_{2r} \rangle'',$$

with $\langle p, q \rangle'' := \langle e_p, I''_{ab} e_q \rangle$. The Weingarten calculus for symplectic groups gives

$$\mathbb{E}[\tilde{w}_{i_1,i_2} \tilde{w}_{i_3,i_4} \cdots \tilde{w}_{i_{2k-1},i_{2k}}] = \sum_{\sigma \in M_{2k}} \Delta'_{\sigma}(\boldsymbol{i}) \sum_{\tau \in M_{2k}} \mathrm{Wg}^{\mathrm{Sp}}(\tau^{-1}\sigma; n) \widetilde{T}(\tau),$$

where

$$\begin{aligned} \widetilde{T}(\tau) :=& \sum_{\boldsymbol{p}=(p_1,\dots,p_{2k})} \Delta'_{\tau}(\boldsymbol{p}) \prod_{r=1}^{k} \langle p_{2r-1}, p_{2r} \rangle'' = \sum_{p_1,\dots,p_{2k}} \prod_{r=1}^{k} \langle p_{\tau(2r-1)}, p_{\tau(2r)} \rangle \langle p_{2r-1}, p_{2r} \rangle'' \\ =& \mathcal{T}_{\tau}(J, JI''_{ab}) = T_{ab}^{\mathrm{C\,II}}(\tau). \end{aligned}$$

□

Example 4.3. $n = a + b$. $\mathrm{Wg}^{\mathrm{C\,II}}(\mathrm{id}_2; a, b) = \frac{a-b}{n}$;

$$\begin{aligned} \mathrm{Wg}^{\mathrm{C\,II}}\left(\left(\begin{smallmatrix} 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \end{smallmatrix}\right); a, b\right) =& \frac{(a-b)^2(2n-1)-n}{n(n-1)(2n+1)}, \\ (-1)\mathrm{Wg}^{\mathrm{C\,II}}\left(\left(\begin{smallmatrix} 1 & 2 & 3 & 4 \\ 1 & 3 & 2 & 4 \end{smallmatrix}\right); a, b\right) = \mathrm{Wg}^{\mathrm{C\,II}}\left(\left(\begin{smallmatrix} 1 & 2 & 3 & 4 \\ 1 & 4 & 2 & 3 \end{smallmatrix}\right); a, b\right) =& \frac{4ab}{n(n-1)(2n+1)}. \end{aligned}$$

Using twisted zonal polynomials Z'_λ (see [Mac, VII, Example 2-7]), we obtain the expansion of $\mathrm{Wg}^{\mathrm{C\,II}}(\cdot; a, b)$ in twisted spherical functions π^λ:

$$\mathrm{Wg}^{\mathrm{C\,II}}(\cdot; a, b) = \frac{2^k k!}{(2k)!} \sum_{\lambda \vdash k} f^{\lambda \cup \lambda} \frac{Z'_\lambda(1^a, (-1)^b)}{Z'_\lambda(1^{a+b})} \pi^\lambda.$$

5 BdG ensembles

In this section, we deal with matrix ensembles of types D III and C I. They are called Bogolioubov-de Gennes (BdG) ensembles.

5.1 Class D III

The setting for D III: $G = \mathrm{O}(2n)$, $K = \mathrm{O}(2n) \cap \mathrm{Sp}(2n) \simeq \mathrm{U}(n)$, $\Omega(g) = (g^{\mathrm{D}})^{-1}$.

Let R be a $2n \times 2n$ Haar orthogonal matrix. We consider the random matrix

$$V = V^{\mathrm{D\,III}} = R^{\mathrm{D}} R,$$

associated to the symmetric space $\mathrm{O}(2n)/\mathrm{U}(n)$.

We define the function $T_n^{\mathrm{D\,III}}$ on S_{2k} by

$$T_n^{\mathrm{D\,III}}(\sigma) = \mathcal{T}_\sigma(I_{2n}, J_n).$$

It satisfies $T_n^{\mathrm{D\,III}}(\zeta\sigma\zeta') = \epsilon(\zeta) T_n^{\mathrm{D\,III}}(\sigma)$ for any $\sigma \in S_{2k}$ and $\zeta, \zeta' \in H_k$. Note that

$$T_n^{\mathrm{D\,III}}(\sigma_\mu) = \prod_{i=1}^{\ell(\mu)} T_n^{\mathrm{D\,III}}(\sigma_{(\mu_i)}) = \prod_{i=1}^{\ell(\mu)} \mathrm{Tr}(J^{\mu_i}) = \begin{cases} (-2n)^{\ell(\mu)} & \text{if } \mu \text{ is even,} \\ 0 & \text{otherwise.} \end{cases}$$

Here a partition μ is said to be *even* if $\mu = 2\nu$ for some partition ν. In particular, $T_n^{\mathrm{D\,III}}(\sigma) = 0$ ($\sigma \in S_{2k}$) if k is odd.

Theorem 5.1. *Let $V = V^{\mathrm{D\,III}}$ be the random matrix defined as above. For a sequence $\boldsymbol{i} = (i_1, i_2, \dots, i_{2k})$ with an even number k, we have*

$$\mathbb{E}[\tilde{v}_{i_1,i_2} \tilde{v}_{i_3,i_4} \cdots \tilde{v}_{i_{2k-1},i_{2k}}] = \sum_{\sigma \in M_{2k}} \Delta_\sigma(\boldsymbol{i}) \mathrm{Wg}^{\mathrm{D\,III}}(\sigma; n).$$

Here $\mathrm{Wg}^{\mathrm{D\,III}}(\cdot;n)$ *is the function on* S_{2k} *defined by*

$$\mathrm{Wg}^{\mathrm{D\,III}}(\cdot;n) = T_n^{\mathrm{D\,III}} \star \mathrm{Wg}^{\mathrm{O}}(\cdot;2n).$$

If k *is odd, then, for any sequence* $\boldsymbol{i} = (i_1, i_2, \dots, i_{2k})$,

$$\mathbb{E}[\tilde{v}_{i_1,i_2}\tilde{v}_{i_3,i_4}\cdots\tilde{v}_{i_{2k-1},i_{2k}}] = 0.$$

Proof. The proof is similar to that of Theorem 4.3. The Weingarten calculus for a Haar orthogonal matrix R gives

$$\mathbb{E}[\tilde{v}_{i_1,i_2}\tilde{v}_{i_3,i_4}\cdots\tilde{v}_{i_{2k-1},i_{2k}}] = \sum_{\sigma\in M_{2k}} \Delta_\sigma(\boldsymbol{i}) \sum_{\tau\in M_{2k}} \mathrm{Wg}^{\mathrm{O}}(\tau^{-1}\sigma;2n)\widetilde{T}(\tau)$$

where

$$\widetilde{T}(\tau) = \sum_{\boldsymbol{p}=(p_1,\dots,p_{2k})} \Delta_\tau(\boldsymbol{p}) \prod_{r=1}^{k} \langle p_{2r-1}, p_{2r}\rangle = \mathcal{T}_\tau(I,J) = T_n^{\mathrm{D\,III}}(\tau)$$

for any $\tau \in S_{2k}$. □

Note that

$$\mathrm{Wg}^{\mathrm{D\,III}}(\zeta\sigma\zeta';n) = \epsilon(\zeta)\mathrm{Wg}^{\mathrm{D\,III}}(\sigma;n) \qquad (\sigma\in S_{2k},\ \zeta,\zeta'\in H_k),$$

and hence $\mathrm{Wg}^{\mathrm{D\,III}}(\zeta;n) = 0$ for $\zeta \in H_k$. Furthermore, $\mathrm{Wg}^{\mathrm{D\,III}}(\sigma;n) = 0$ ($\sigma \in S_{2k}$) if k is odd.

Example 5.1. Since $\left(\begin{smallmatrix}1&2&3&4\\1&3&2&4\end{smallmatrix}\right) = (3\ 4)\left(\begin{smallmatrix}1&2&3&4\\1&4&2&3\end{smallmatrix}\right)$ and $(3\ 4) \in H_2$, we have

$$(-1)\mathrm{Wg}^{\mathrm{D\,III}}\left(\left(\begin{smallmatrix}1&2&3&4\\1&3&2&4\end{smallmatrix}\right);n\right) = \mathrm{Wg}^{\mathrm{D\,III}}\left(\left(\begin{smallmatrix}1&2&3&4\\1&4&2&3\end{smallmatrix}\right);n\right) = \frac{-1}{2n-1}.$$

5.2 Class C I

The setting for C I: $G = \mathrm{Sp}(2n)$, $K = \mathrm{U}(n)$, $\Omega(g) = I'_{nn} g I'_{nn}$. Here the fix-point set of Ω in G is

$$\left\{ \begin{pmatrix} U & O \\ O & \overline{U} \end{pmatrix} \ \middle|\ U \in \mathrm{U}(n) \right\} \simeq \mathrm{U}(n).$$

Let S be a $2n \times 2n$ Haar symplectic matrix. We consider the random matrix

$$W^{\mathrm{C\,I}} = S^{\mathrm{D}} I'_{nn} S,$$

instead of $V^{\mathrm{C\,I}} = \Omega(S)^{-1}S = I'_{nn}S^{D}I'_{nn}S$, which is associated to the symmetric space $\mathrm{Sp}(2n)/\mathrm{U}(n)$.

We define the function $T_n^{\mathrm{C\,I}}$ on S_{2k} by

$$T_n^{\mathrm{C\,I}}(\sigma) = \mathcal{T}_\sigma(J, JI'_{nn}).$$

Since $JI'_{nn} = -\left(\begin{smallmatrix} O_n & I_n \\ I_n & O_n \end{smallmatrix}\right)$ is symmetric and J is skew-symmetric, the function $T_n^{\mathrm{C\,I}}$ satisfies $T_n^{\mathrm{C\,I}}(\zeta\sigma\zeta') = T_n^{\mathrm{C\,I}}(\sigma)\epsilon(\zeta')$ for $\sigma \in S_{2k}$ and $\zeta, \zeta' \in H_k$. For $\sigma = \sigma_\mu$, we have

$$\begin{aligned} T_n^{\mathrm{C\,I}}(\sigma_\mu) &= \prod_{i=1}^{\ell(\mu)} T_n^{\mathrm{C\,I}}(\sigma_{(\mu_i)}) = \prod_{i=1}^{\ell(\mu)} [-\operatorname{Tr}(JJI'_{nn})^{\mu_i}] = (-1)^{\ell(\mu)} \prod_{i=1}^{\ell(\mu)} \operatorname{Tr}(-I'_{nn})^{\mu_i} \\ &= \begin{cases} (-2n)^{\ell(\mu)} & \text{if } \mu \text{ is even,} \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

In particular, $T_n^{\mathrm{C\,I}}(\sigma) = T_n^{\mathrm{D\,III}}(\sigma^{-1}) = \pm T_n^{\mathrm{D\,III}}(\sigma)$ for all $\sigma \in S_{2k}$.

Theorem 5.2. *Let $W = W^{\mathrm{C\,I}}$ be the random matrix defined as above. For a sequence $\boldsymbol{i} = (i_1, i_2, \dots, i_{2k})$ with an even number k, we have*

$$\mathbb{E}[\tilde{w}_{i_1,i_2}\tilde{w}_{i_3,i_4}\cdots\tilde{w}_{i_{2k-1},i_{2k}}] = \sum_{\sigma \in M_{2k}} \Delta'_\sigma(\boldsymbol{i}) \mathrm{Wg}^{\mathrm{C\,I}}(\sigma; n).$$

Here $\mathrm{Wg}^{\mathrm{C\,I}}(\cdot; n)$ is the function on S_{2k} defined by

$$\mathrm{Wg}^{\mathrm{C\,I}}(\cdot; n) = T_n^{\mathrm{C\,I}} \star \mathrm{Wg}^{\mathrm{Sp}}(\cdot; n).$$

If k is odd, then, for any sequence $\boldsymbol{i} = (i_1, i_2, \dots, i_{2k})$,

$$\mathbb{E}[\tilde{w}_{i_1,i_2}\tilde{w}_{i_3,i_4}\cdots\tilde{w}_{i_{2k-1},i_{2k}}] = 0.$$

Proof. It is proved in a usual way. The Weingarten calculus for a Haar symplectic matrix S gives

$$\mathbb{E}[\tilde{w}_{i_1,i_2}\tilde{w}_{i_3,i_4}\cdots\tilde{w}_{i_{2k-1},i_{2k}}] = \sum_{\sigma \in M_{2k}} \Delta'_\sigma(\boldsymbol{i}) \sum_{\tau \in M_{2k}} \mathrm{Wg}^{\mathrm{Sp}}(\tau^{-1}\sigma; n)\widetilde{T}(\tau)$$

where

$$\widetilde{T}(\tau) := \sum_{\boldsymbol{p}=(p_1,\dots,p_{2k})} \Delta'_\tau(\boldsymbol{p}) \prod_{r=1}^{k} \langle e_{p_{2r-1}}, I'_{nn} e_{p_{2r}} \rangle = \mathcal{T}_\tau(J, JI'_{nn}) = T_n^{\mathrm{C\,I}}(\tau)$$

for any $\tau \in S_{2k}$. □

The function $\mathrm{Wg}^{\mathrm{C\,I}}(\cdot; n)$ on S_{2k} satisfies

$$\mathrm{Wg}^{\mathrm{C\,I}}(\zeta\sigma\zeta'; n) = \epsilon(\zeta')\mathrm{Wg}^{\mathrm{C\,I}}(\sigma; n) \qquad (\sigma \in S_{2k},\ \zeta, \zeta' \in H_k),$$

and hence $\mathrm{Wg}^{\mathrm{C\,I}}(\zeta; n) = 0$ for $\zeta \in H_k$. Furthermore, $\mathrm{Wg}^{\mathrm{C\,I}}(\sigma; n) = 0$ ($\sigma \in S_{2k}$) if k is odd.

Example 5.2.

$$\mathrm{Wg}^{\mathrm{C\,I}}\left(\left(\begin{smallmatrix} 1 & 2 & 3 & 4 \\ 1 & 3 & 2 & 4 \end{smallmatrix}\right); n\right) = \mathrm{Wg}^{\mathrm{C\,I}}\left(\left(\begin{smallmatrix} 1 & 2 & 3 & 4 \\ 1 & 4 & 2 & 3 \end{smallmatrix}\right); n\right) = \frac{-1}{2n+1}.$$

6 Conclusion

We have made methods for computations of moments of matrix elements from classical compact Lie groups and classical compact symmetric spaces. Write $\mathcal{C} =$ A, B/D, C for unitary, orthogonal, symplectic groups, respectively. The moment for a classical group is given by the double sum

$$\sum_{\sigma}\sum_{\tau}(\Delta\text{-function in }\sigma)\times(\Delta\text{-function in }\tau)\times \mathrm{Wg}^{\mathcal{C}}(\sigma^{-1}\tau;n),$$

whereas that for a classical compact symmetric space is given by the single sum

$$\sum_{\sigma}(\Delta\text{-function in }\sigma)\times \mathrm{Wg}^{\mathcal{C}}(\sigma;n).$$

Here the Δ-function is

- $\delta_\sigma(\cdot,\cdot)$ defined in (2.2) if $\mathcal{C} =$ A, A I, A II, A III,
- $\Delta_\sigma(\cdot)$ defined in (2.6) if $\mathcal{C} =$ B/D, BD I, D III,
- $\Delta'_\sigma(\cdot)$ defined in (2.12) if $\mathcal{C} =$ C, C I, C II,

and the Weingarten function $\mathrm{Wg}^{\mathcal{C}}$ belongs to

- $\mathcal{Z}(L(S_k))$ if $\mathcal{C} =$ A, A III,
- $L(S_{2k}, H_k)$ if $\mathcal{C} =$ B/D, A I, BD I,
- $L^{\epsilon}(S_{2k}, H_k)$ if $\mathcal{C} =$ C, A II, C II.

The Weingarten functions for $\mathcal{C} =$ D III or C I differ from others. In fact, if we let $F(\sigma)$ to be $\mathrm{Wg}^{\mathrm{D\,III}}(\sigma;n)$ or $\mathrm{Wg}^{\mathrm{C\,I}}(\sigma^{-1};n)$, then F is the function on S_{2k} satisfying the property

$$F(\zeta\sigma\zeta') = \epsilon(\zeta)F(\sigma) \qquad (\sigma \in S_{2k},\ \zeta,\zeta' \in H_k).$$

In particular, F identically vanishes on H_k, and, moreover, on S_{2k} if k is odd. Weingarten functions except D III and C I have Fourier expansions in χ^λ, ω^λ, or π^λ. However, we could not find such expansions for these two cases.

Acknowledgments

The author's work was supported by JSPS Grant-in-Aid for Young Scientists (B) 22740060.

References

[Ca] E. Cartan, Sur certaines formes Riemanniennes remarquables des géométries à groupe fondamental simple, (French), Ann. Sci. École Norm. Sup. (3) **44** (1927), 345–467.

[CaM] M. Caselle and U. Magnea, Random matrix theory and symmetric spaces, Physics Reports **394** (2004), 41–156.

[Co] B. Collins, Moments and cumulants of polynomial random variables on unitary groups, the Itzykson-Zuber integral, and free probability, Int. Math. Res. Not. (2003), no. 17, 953–982.

[CFN] B. Collins, M. Fukuda, and I. Nechita, Towards a state minimizing the output entropy of a tensor product of random quantum channels, J. Math. Phys. **53** (2012), no. 3, 032203, 20 pp.

[CM] B. Collins and S. Matsumoto, On some properties of orthogonal Weingarten functions, J. Mathematical Phys. **50** (2009), no. 11, 113516, 14 pp.

[CMS] B. Collins, S. Matsumoto, and N. Saad, Integration of invariant matrices and application to statistics, arXiv:1205.0956, 19 pp.

[CN] B. Collins and I. Nechita, Gaussianization and eigenvalue statistics for random quantum channels (III), Ann. Appl. Probab. **21** (2011), no. 3, 1136–1179.

[CSn] B. Collins and P. Śniady, Integration with respect to the Haar measure on unitary, orthogonal and symplectic group, Comm. Math. Phys. **264** (2006), no. 3, 773–795.

[CSt] B. Collins and M. Stolz, Borel theorems for random matrices from the classical compact symmetric spaces, The Annals of Probability **36** (2008), no. 3, 876–895.

[Da] A. Dahlqvist, Integration formulas for Brownian motion on classical groups, arXiv:1212.5107v1.

[Du] E. Dueñez, Random matrix ensembles associated to compact symmetric spaces, Commun. Math. Phys. **244** (2004), 29–61.

[Dy] F. J. Dyson, Statistical theory of the energy levels of complex systems. I, II, III, Math. Phys. **3** (1962), 140–156, 157–165, 166–175.

[F] P. J. Forrester, Log-gases and random matrices, London Mathematical Society Monographs Series **34**, Princeton University Press, 2010.

[GGJ] I. P. Goulden, M. Guay-Paquet, and J. Novak, Monotone Hurwitz numbers and the HCIZ integral, arXiv:1107.1015v3.

[GW] R. Goodman and N. R. Wallach, Symmetry, representations, and invariants, Graduate Texts in Mathematics **255**, Springer, Dordrecht, 2009.

[Mac] I. G. Macdonald, Symmetric Functions and Hall Polynomials, second ed., Oxford University Press, Oxford, 1995.

[M1] S. Matsumoto, Jucys-Murphy elements, orthogonal matrix integrals, and Jack measures, The Ramanujan J. **26** (2011), no. 1, 69–107.

[M2] S. Matsumoto, General moments of matrix elements from circular orthogonal ensembles, Random Matrices: Theory and Applications **1** (2012), no. 3, 125005, 18 pp.

[M3] S. Matsumoto, Moments of a single entry of circular orthogonal ensembles and Weingarten calculus, Let. Math. Phys. **103** (2013), 113–130.

[MN] S. Matsumoto and J. Novak, Jucys-Murphy elements and unitary matrix integrals, International Mathematics Research Notices (2013), no. 2, 362–397.

[N] J. Novak, Jucys-Murphy elements and the unitary Weingarten function, Banach Center Publications **89** (2010), 231–235.

[S] A. J. Scott, Optimizing quantum process tomography with unitary 2-designs, J. Phys. A: Math. Theor. **41** (2008), 055308, 26pp.

[W] D. Weingarten, Asymptotic behavior of group integrals in the limit of infinite rank, J. Mathematical Phys. **19** (1978), no. 5, 999–1001.

[Z] P. Zinn-Justin, Jucys-Murphy elements and Weingarten matrices, Let. Math. Phys. **91** (2010), 119–127.

SHO MATSUMOTO
Graduate School of Mathematics, Nagoya University, Nagoya, 464-8602, Japan.
`sho-matsumoto@math.nagoya-u.ac.jp`